

10/783,801

Connecting via Winsock to STN

Welcome to STN International! Enter x:x :

LOGINID:sssptal201txs

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page URLs for STN Seminar Schedule - N. America
NEWS	2		"Ask CAS" for self-help around the clock
NEWS	3	JAN 17	Pre-1988 INPI data added to MARPAT
NEWS	4	FEB 21	STN AnaVist, Version 1.1, lets you share your STN AnaVist visualization results
NEWS	5	FEB 22	The IPC thesaurus added to additional patent databases on STN
NEWS	6	FEB 22	Updates in EPFULL; IPC 8 enhancements added
NEWS	7	FEB 27	New STN AnaVist pricing effective March 1, 2006
NEWS	8	MAR 03	Updates in PATDPA; addition of IPC 8 data without attributes
NEWS	9	MAR 22	EMBASE is now updated on a daily basis
NEWS	10	APR 03	New IPC 8 fields and IPC thesaurus added to PATDPAFULL
NEWS	11	APR 03	Bibliographic data updates resume; new IPC 8 fields and IPC thesaurus added in PCTFULL
NEWS	12	APR 04	STN AnaVist \$500 visualization usage credit offered
NEWS	13	APR 12	LINSPEC, learning database for INSPEC, reloaded and enhanced
NEWS	14	APR 12	Improved structure highlighting in FQHIT and QHIT display in MARPAT
NEWS	15	APR 12	Derwent World Patents Index to be reloaded and enhanced during second quarter; strategies may be affected
NEWS	16	MAY 10	CA/CAPLUS enhanced with 1900-1906 U.S. patent records
NEWS	17	MAY 11	KOREAPAT updates resume
NEWS	18	MAY 19	Derwent World Patents Index to be reloaded and enhanced
NEWS	19	MAY 30	IPC 8 Rolled-up Core codes added to CA/CAPLUS and USPATFULL/USPAT2
NEWS	20	MAY 30	The F-Term thesaurus is now available in CA/CAPLUS
NEWS	21	JUN 02	The first reclassification of IPC codes now complete in INPADOC
NEWS EXPRESS			FEBRUARY 15 CURRENT VERSION FOR WINDOWS IS V8.01a, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 19 DECEMBER 2005. V8.0 AND V8.01 USERS CAN OBTAIN THE UPGRADE TO V8.01a AT http://download.cas.org/express/v8.0-Discover/
NEWS HOURS			STN Operating Hours Plus Help Desk Availability
NEWS LOGIN			Welcome Banner and News Items
NEWS IPC8			For general information regarding STN implementation of IPC 8
NEWS X25			X.25 communication option no longer available after June 2006

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific

10/783,801

research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 09:38:37 ON 07 JUN 2006

=> file reg

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.21	0.21

FULL ESTIMATED COST

FILE 'REGISTRY' ENTERED AT 09:38:50 ON 07 JUN 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 6 JUN 2006 HIGHEST RN 887000-62-6

DICTIONARY FILE UPDATES: 6 JUN 2006 HIGHEST RN 887000-62-6

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 6, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*

Structure search iteration limits have been increased. See HELP SLIMITS for details.


REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/ONLINE/UG/regprops.html>

=>

Uploading C:\Program Files\Stnexp\Queries\107838013.str

10/783,801



```
chain nodes :
17 18 19 20 21
ring nodes :
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16
chain bonds :
1-18 3-17 8-12 9-19 15-20 16-21
ring bonds :
1-2 1-6 2-3 3-4 4-5 5-6 5-7 6-10 7-8 8-9 9-10 11-12 11-16 12-13 13-14
14-15 15-16
exact/norm bonds :
1-18 3-17 5-7 6-10 7-8 8-9 9-10 9-19 15-20 16-21
exact bonds :
8-12
normalized bonds :
1-2 1-6 2-3 3-4 4-5 5-6 11-12 11-16 12-13 13-14 14-15 15-16
```

```
Match level :
1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:CLASS 18:CLASS 19:CLASS
20:CLASS 21:CLASS
```

L1 STRUCTURE UPLOADED

=> s 11

SAMPLE SEARCH INITIATED 09:39:16 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 1036 TO ITERATE

100.0% PROCESSED 1036 ITERATIONS 50 ANSWERS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**

10/783,801

BATCH **COMPLETE**
PROJECTED ITERATIONS: 18789 TO 22651
PROJECTED ANSWERS: 5639 TO 7841

L2 50 SEA SSS SAM L1

=> s l1 ful
FULL SEARCH INITIATED 09:39:24 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 20139 TO ITERATE

100.0% PROCESSED 20139 ITERATIONS 6505 ANSWERS
SEARCH TIME: 00.00.01

L3 6505 SEA SSS FUL L1

=> file caplus
COST IN U.S. DOLLARS SINCE FILE TOTAL
ENTRY SESSION
FULL ESTIMATED COST 166.94 167.15

FILE 'CAPLUS' ENTERED AT 09:39:34 ON 07 JUN 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 7 Jun 2006 VOL 144 ISS 24
FILE LAST UPDATED: 6 Jun 2006 (20060606/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s l3
L4 16900 L3

=> s l4 and (epicatechin or catechin)
4775 EPICATECHIN
63 EPICATECHINS
4790 EPICATECHIN
(EPICATECHIN OR EPICATECHINS)
8467 CATECHIN
3106 CATECHINS
9554 CATECHIN
(CATECHIN OR CATECHINS)
L5 8190 L4 AND (EPICATECHIN OR CATECHIN)

=> s l5 and dimer
103875 DIMER

10/783,801

59006 DIMERS
136839 DIMER
(DIMER OR DIMERS)

L6 408 L5 AND DIMER

=> s 16 and (4,8)

5339334 4
2704086 8
112120 4,8
(4(W)8)

L7 18 L6 AND (4,8)

=> s 15 and (process or make or made or prepar? or synthes? or method)

2251291 PROCESS
1525430 PROCESSES
3360296 PROCESS
(PROCESS OR PROCESSES)

231644 MAKE
180294 MAKES
399557 MAKE
(MAKE OR MAKES)

1207430 MADE
25 MADES
1207451 MADE
(MADE OR MADES)

1647869 PREPAR?
122650 PREP
2150 PREPS
124594 PREP
(PREP OR PREPS)

2008322 PREPD
17 PREPDS
2008334 PREPD
(PREPD OR PREPDS)

121223 PREPG
12 PREPGS
121234 PREPG
(PREPG OR PREPGS)

2711598 PREPN
204470 PREPNS
2865843 PREPN
(PREPN OR PREPNS)

4743234 PREPAR?
(PREPAR? OR PREP OR PREPD OR PREPG OR PREPN)

1535222 SYNTHES?

3100220 METHOD

1272104 METHODS

4013969 METHOD
(METHOD OR METHODS)

L8 3366 L5 AND (PROCESS OR MAKE OR MADE OR PREPAR? OR SYNTHES? OR METHOD
)

=> s 16 and (process or make or made or prepar? or synthes? or method)

2251291 PROCESS
1525430 PROCESSES
3360296 PROCESS
(PROCESS OR PROCESSES)

231644 MAKE
180294 MAKES
399557 MAKE

10/783,801

(MAKE OR MAKES)
1207430 MADE
25 MADES
1207451 MADE
(MADE OR MADES)
1647869 PREPAR?
122650 PREP
2150 PREPS
124594 PREP
(PREP OR PREPS)
2008322 PREPD
17 PREPDS
2008334 PREPD
(PREPD OR PREPDS)
121223 PREPG
12 PREPGS
121234 PREPG
(PREPG OR PREPGS)
2711598 PREPN
204470 PREPNS
2865843 PREPN
(PREPN OR PREPNS)
4743234 PREPAR?
(PREPAR? OR PREP OR PREPD OR PREPG OR PREPN)
1535222 SYNTHES?
3100220 METHOD
1272104 METHODS
4013969 METHOD
(METHOD OR METHODS)

L9 185 L6 AND (PROCESS OR MAKE OR MADE OR PREPAR? OR SYNTHES? OR METHOD
)

=> s l9 and (oxidation or oxidi?)
433563 OXIDATION
4850 OXIDATIONS
434826 OXIDATION
(OXIDATION OR OXIDATIONS)
742953 OXIDN
9337 OXIDNS
744911 OXIDN
(OXIDN OR OXIDNS)
880077 OXIDATION
(OXIDATION OR OXIDN)
404853 OXIDI?

L10 27 L9 AND (OXIDATION OR OXIDI?)

=> d l10 ibib hitstr abs 1-27

L10 ANSWER 1 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2005:1273656 CAPLUS

DOCUMENT NUMBER: 144:308365

TITLE: Transparent testal0 encodes a laccase-like enzyme
involved in oxidative polymerization of flavonoids in
Arabidopsis seed coat


AUTHOR(S): Pourcel, Lucille; Routaboul, Jean-Marc; Kerhoas,
Lucien; Caboche, Michel; Lepiniec, Loic; Debeaujon,
Isabelle

CORPORATE SOURCE: Laboratoire de Biologie des Semences, Unite Mixte de
Recherche 204, Institut National de la Recherche
Agronomique/Institut National Agronomique

10/783,801

Paris-Grignon, Institut Jean-Pierre Bourgin, Institut
National de la Recherche Agronomique/Institut National
Agronomique Paris-Grignon, Versailles, 78026, Fr.
SOURCE: Plant Cell (2005), 17(11), 2966-2980
CODEN: PLCEEW; ISSN: 1040-4651
PUBLISHER: American Society of Plant Biologists
DOCUMENT TYPE: Journal
LANGUAGE: English
IT **490-46-0, Epicatechin**
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(transparent testa10 encodes laccase-like enzyme involved in oxidative
polymerization of flavonoids in Arabidopsis seed coat)
RN 490-46-0 CAPLUS
CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-,
(2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




AB The Arabidopsis thaliana transparent testa10 (tt10) mutant exhibits a delay in developmentally determined browning of the seed coat, also called the testa. Seed coat browning is caused by the **oxidation** of flavonoids, particularly proanthocyanidins, which are polymers of flavan-3-ol subunits such as **epicatechin** and **catechin**. The tt10 mutant seeds accumulate more **epicatechin** monomers and more soluble proanthocyanidins than wild-type seeds. Moreover, intact testa cells of tt10 cannot trigger H2O2-independent browning in the presence of **epicatechin** and **catechin**, in contrast with wild-type cells. UV-visible light detection and mass spectrometry revealed that the major **oxidation** products obtained with **epicatechin** alone are yellow **dimers** called dehydrodiepicatechin A. These products differ from proanthocyanidins in the nature and position of their interflavan linkages. Flavonol composition was also affected in tt10 seeds, which exhibited a higher ratio of quercetin rhamnoside monomers vs. **dimers** than wild-type seeds. We identified the TT10 gene by a candidate gene approach. TT10 encodes a protein with strong similarity to laccase-like polyphenol oxidases. It is expressed essentially in developing testa, where it colocalizes with the flavonoid end products proanthocyanidins and flavonols. Together, these data establish that TT10 is involved in the oxidative polymerization of flavonoids and functions as a laccase-type flavonoid oxidase.
REFERENCE COUNT: 57 THERE ARE 57 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 2 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2005:1165929 CAPLUS
DOCUMENT NUMBER: 144:16442

10/783,801


TITLE: Phenolics as potential antioxidant therapeutic agents:
Mechanism and actions
AUTHOR(S): Soobrattee, M. A.; Neergheen, V. S.; Luximon-Ramma,
A.; Aruoma, O. I.; Bahorun, T.
CORPORATE SOURCE: Department of Biosciences, Faculty of Science,
University of Mauritius, Reduit, Mauritius
SOURCE: Mutation Research (2005), 579(1-2), 200-213
CODEN: MUREAV; ISSN: 0027-5107
PUBLISHER: Elsevier B.V.
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 154-23-4, (+)-Catechin 490-46-0, (-)-
Epicatechin 528-58-5, Cyanidin chloride 970-74-1
, (-)-Epigallocatechin 989-51-5, (-)-Epigallocatechin gallate
1257-08-5 20315-25-7, Procyanidin B1 29106-49-8
, Procyanidin B2
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL
(Biological study); USES (Uses)
(phenolics as potential antioxidant therapeutic agents and mechanism
and actions)
RN 154-23-4 CAPLUS
CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-,
(2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



RN 490-46-0 CAPLUS
CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-,
(2R,3R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



10/783,801

RN 528-58-5 CAPLUS

CN 1-Benzopyrylium, 2-(3,4-dihydroxyphenyl)-3,5,7-trihydroxy-, chloride (9CI)
(CA INDEX NAME)




● Cl⁻

RN 970-74-1 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 3,4-dihydro-2-(3,4,5-trihydroxyphenyl)-,
(2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




RN 989-51-5 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, (2R,3R)-3,4-dihydro-5,7-dihydroxy-2-
(3,4,5-trihydroxyphenyl)-2H-1-benzopyran-3-yl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).


10/783,801



RN 1257-08-5 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, (2R,3R)-2-(3,4-dihydroxyphenyl)-3,4-dihydro-5,7-dihydroxy-2H-1-benzopyran-3-yl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




RN 20315-25-7 CAPLUS

CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3R,3'S,4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/783,801



RN 29106-49-8 CAPLUS
CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3R,3'R,4R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



AB Accumulating chemical, biochem., clin. and epidemiol. evidence supports the chemoprotective effects of phenolic antioxidants against oxidative stress-mediated disorders. The pharmacol. actions of phenolic antioxidants stem mainly from their free radical scavenging and metal chelating properties as well as their effects on cell signaling pathways and on gene expression. The antioxidant capacities of phenolic compds. that are widely distributed in plant-based diets were assessed by the Trolox equivalent antioxidant capacity (TEAC), the ferric reducing antioxidant

power (FRAP), the hypochlorite scavenging capacity, the deoxyribose **method** and the copper-phenanthroline-dependent DNA **oxidn** . assays. Based on the TEAC, FRAP and hypochlorite scavenging data, the observed activity order was: procyanidin **dimer** > flavanol > flavanol > hydroxycinnamic acids > simple phenolic acids. Among the flavanol aglycons, the antioxidant propensities decrease in the order quercetin, myricetin and kaempferol. Gallic acid and rosmarinic acid were the most potent antioxidants among the simple phenolic and hydroxycinnamic acids, resp. Ferulic acid displayed the highest inhibitory activity against deoxyribose degradation but no structure-activity relation could be established for the activities of the phenolic compds. in the deoxyribose assay. The efficacies of the phenolic compds. differ depending on the mechanism of antioxidant action in the resp. assay used, with procyanidin **dimers** and flavan-3-ols showing very potent activities in most of the systems tested. Compared to the physiol. active (glutathione, α -tocopherol, ergothioneine) and synthetic (Trolox, BHA, BHT) antioxidants, these compds. exhibited much higher efficacy. Plant-derived phenolics represents good sources of natural antioxidants, however, further investigation on the mol. mechanism of action of these phytochems. is crucial to the evaluation of their potential as prophylactic agents.

REFERENCE COUNT: 55 THERE ARE 55 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 3 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2005:904441 CAPLUS

DOCUMENT NUMBER: 143:404966

TITLE: A novel black tea pigment and two new **oxidation** products of epigallocatechin-3-O-gallate

AUTHOR(S): Tanaka, Takashi; Matsuo, Yosuke; Kouno, Isao
CORPORATE SOURCE: Graduate School of Biomedical Sciences, Nagasaki University, Nagasaki, 852-8521, Japan

SOURCE: Journal of Agricultural and Food Chemistry (2005), 53(19), 7571-7578

CODEN: JAFCAU; ISSN: 0021-8561

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 989-51-5, Epigallocatechin-3-O-gallate 4670-05-7D,

Theaflavin, derivs. 89064-31-3, Theasinensin A

867278-31-7 867358-20-1, Dehydrotheasinensin AQ


RL: BSU (Biological study, unclassified); BIOL (Biological study)
(black tea pigment and two new **oxidation** products of epigallocatechin-3-O-gallate)

RN 989-51-5 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, (2R,3R)-3,4-dihydro-5,7-dihydroxy-2-(3,4,5-trihydroxyphenyl)-2H-1-benzopyran-3-yl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).


10/783,801



RN 4670-05-7 CAPLUS

CN 5H-Benzocyclohepten-5-one, 1,8-bis[(2R,3R)-3,4-dihydro-3,5,7-trihydroxy-2H-1-benzopyran-2-yl]-3,4,6-trihydroxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).




RN 89064-31-3 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, [(1R)-4,4',5,5',6,6'-hexahydroxy[1,1'-biphenyl]-2,2'-diyl]bis[(2R,3R)-3,4-dihydro-5,7-dihydroxy-2H-1-benzopyran-2,3-diyl] ester (9CI) (CA INDEX NAME)

PAGE 1-A





PAGE 2-A



RN 867278-31-7 CAPLUS

CN [1,1'-Biphenyl]-2-carboxylic acid, 2'-[(2R,3R)-3,4-dihydro-5,7-dihydroxy-3-(3,4,5-trihydroxyphenoxy)-2H-1-benzopyran-2-yl]-3',4,4',5,5',6-hexahydroxy-, (2R,3R)-2-[6'-[(2R,3R)-3,4-dihydro-5,7-dihydroxy-3-[(3,4,5-trihydroxybenzoyl)oxy]-2H-1-benzopyran-2-yl]-2',3',4,4',5,6-hexahydroxy[1,1'-biphenyl]-2-yl]-3,4-dihydro-5,7-dihydroxy-2H-1-benzopyran-3-yl ester (9CI) (CA INDEX NAME)



Absolute stereochemistry.



RN 867358-20-1 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, (7,8-dihydro-3,4,9-trihydroxy-7,8-dioxo-1,5a(6H)-dibenzofurandiyl)bis(3,4-dihydro-5,7-dihydroxy-2H-1-benzopyran-2,3-diyl) ester, stereoisomer (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 625371-09-7, Dehydrotheasinensin A


RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)

(black tea pigment and two new **oxidation** products of epigallocatechin-3-O-gallate)

RN 625371-09-7 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, [(2R,6S)-3,6-dihydro-2,9,10,11,11-pentahydroxy-3-oxo-2,6-methano-2H-1-benzoxocin-5,7-diyl]bis[(2R,3R)-3,4-dihydro-5,7-dihydroxy-2H-1-benzopyran-2,3-diyl] ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



AB During tea fermentation, **oxidation**-reduction dismutation of a number of quinone metabolites of tea **catechins** yields numerous minor products, which **make** it difficult to sep. and purify black tea polyphenols. In this study, epigallocatechin-3-O-gallate was enzymically **oxidized** and then the unstable quinone metabolites in the **oxidation** mixture were hydrogenated with 2-mercaptoethanol to reduce production of inseparable minor dismutation products. As a result, three new **oxidation** products including a new black tea pigment were isolated, and their structures were determined based on chemical and spectroscopic data. Dehydrotheasinensin AQ is a new reddish-orange pigment with a 1,2-diketone structure, and its presence in com. black tea was confirmed. In addition, a new quinone **dimer** with a complex caged structure and a trimer of epigallocatechin-3-O-gallate were also isolated and their production mechanisms are proposed. The presence of this trimer suggested participation of galloyl quinones in production of minor polyphenols in black tea.

REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 4 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2004:513695 CAPLUS

DOCUMENT NUMBER: 141:410544

TITLE: Study of the **oxidation processes** of **catechins** by on-line electrolysis/ESI/MS

AUTHOR(S): Yamaguchi, Masashi; Mizooku, Yasuo; Osakai, Toshiyuki; Kimoto, Takashi; Arakawa, Ryuichi

CORPORATE SOURCE: Department of Applied Chemistry, Faculty of Engineering, Kansai University, Suita-shi, Osaka, 564-8680, Japan

SOURCE: Bunseki Kagaku (2004), 53(6), 547-553
CODEN: BNSKAK; ISSN: 0525-1931

PUBLISHER: Nippon Bunseki Kagakkai

DOCUMENT TYPE: Journal

LANGUAGE: Japanese

IT 791816-85-8


10/783,801

RL: CPS (Chemical process); FMU (Formation, unclassified); PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); FORM (Formation, nonpreparative); PROC (Process); RACT (Reactant or reagent)
(oxidation of catechins examined by online electrolysis/ESI/MS)

RN 791816-85-8 CAPLUS

CN 3,5-Cyclohexadiene-1,2-dione, 3-hydroxy-5-[(2R,2'R,3R,3'R)-3,3',4,4'-tetrahydro-3,3',5,5',7,7'-hexahydroxy-2'-(3,4,5-trihydroxyphenyl)[4,8'-bi-2H-1-benzopyran]-2-yl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 970-74-1, (-)-Epigallocatechin 989-51-5,

(-)-Epigallocatechin gallate 1257-08-5


RL: CPS (Chemical process); FMU (Formation, unclassified); PEP (Physical, engineering or chemical process); RCT (Reactant); FORM (Formation, nonpreparative); PROC (Process); RACT (Reactant or reagent)

(oxidation of catechins examined by online electrolysis/ESI/MS)

RN 970-74-1 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 3,4-dihydro-2-(3,4,5-trihydroxyphenyl)-, (2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




10/783,801

RN 989-51-5 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, (2R,3R)-3,4-dihydro-5,7-dihydroxy-2-(3,4,5-trihydroxyphenyl)-2H-1-benzopyran-3-yl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 1257-08-5 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, (2R,3R)-2-(3,4-dihydroxyphenyl)-3,4-dihydro-5,7-dihydroxy-2H-1-benzopyran-3-yl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



AB Much attention was focused on the strong antioxidant activity and antitumorigenic property of **catechins** extracted from green tea. However, not many studies on the **oxidation process** of **catechins** were conducted. The authors employed an online electrochem./electrospray ionization mass spectrometry (EC/ESI-MS) technique to identify unstable **oxidation** products of **catechins**. The authors have succeeded in detecting **dimer**

products formed by the electrolysis of (-)-epigallocatechin, (-)-**epicatechin** gallate and (-)-epigallocatechin gallate. EC/ESI-MS was a useful technique to study electrochem. **oxidation process** of antioxidants.

L10 ANSWER 5 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2003:690290 CAPLUS

DOCUMENT NUMBER: 139:337052

TITLE: **Oxidation of tea catechins:**
chemical structures and reaction mechanism

AUTHOR(S): Tanaka, Takashi; Kouno, Isao

CORPORATE SOURCE: Department of Molecular Medicinal Sciences, Graduate School of Biomedical Sciences, Nagasaki University, Nagasaki, 852-8521, Japan

SOURCE: Food Science and Technology Research (2003), 9(2), 128-133

CODEN: FSTRFS; ISSN: 1344-6606

PUBLISHER: Japanese Society for Food Science and Technology

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English


IT **4670-05-7**, Theaflavin

RL: BCP (Biochemical process); BIOL (Biological study); PROC (Process)
(**oxidation of tea catechins** related to polyphenols in black tea)

RN 4670-05-7 CAPLUS

CN 5H-Benzocyclohepten-5-one, 1,8-bis[(2R,3R)-3,4-dihydro-3,5,7-trihydroxy-2H-1-benzopyran-2-yl]-3,4,6-trihydroxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



AB A review. Black tea accounts for almost 80% of the world's tea production and is the most important source of polyphenol in the world. However, little has been known about the chemical of black tea polyphenols due to their complexity. Since most of the black tea polyphenols are produced by enzymic **oxidation** of green tea **catechins**, in vitro model fermentation expts. using purified **catechins** are very useful, and recently structures of some novel **oxidation** products of theaflavins, black tea pigments, have been elucidated. In addition, accumulation of unstable **dimer** quinones of epigallocatechin and its gallate during tea fermentation has been demonstrated, and the **dimer** quinones are converted to theasinensins, another major polyphenol characteristic of black tea, on heating. Formation and degradation of theaflavins and epigallocatechin **dimer** quinones are major pathways in **catechin oxidation** during tea fermentation and

10/783,801

understanding the chemical mechanism is important in clarifying black tea polyphenols.

REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 6 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2003:613003 CAPLUS

DOCUMENT NUMBER: 139:393356

TITLE: **Oxidation** mechanism of **catechins** and related polyphenols

AUTHOR(S): Tanaka, Takashi; Mine, Chie; Matsuda, Miyuki; Inoue, Kyoko; Kouno, Isao

CORPORATE SOURCE: School of Pharmaceutical Sciences, Nagasaki University, Japan

SOURCE: Tennen Yuki Kagobutsu Toronkai Koen Yoshishu (2001), 43rd, 329-334

CODEN: TYKYDS

PUBLISHER: Nippon Kagakkai

DOCUMENT TYPE: Journal; General Review

LANGUAGE: Japanese

IT 4670-05-7, Theaflavin 89064-31-3, Theasinensin A


RL: BSU (Biological study, unclassified); BIOL (Biological study)

(**oxidation** mechanism of **catechins** and related polyphenols)

RN 4670-05-7 CAPLUS

CN 5H-Benzocyclohepten-5-one, 1,8-bis[(2R,3R)-3,4-dihydro-3,5,7-trihydroxy-2H-1-benzopyran-2-yl]-3,4,6-trihydroxy- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).




RN 89064-31-3 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, [(1R)-4,4',5,5',6,6'-hexahydroxy[1,1'-biphenyl]-2,2'-diyl]bis[(2R,3R)-3,4-dihydro-5,7-dihydroxy-2H-1-benzopyran-2,3-diyl] ester (9CI) (CA INDEX NAME)

PAGE 1-A



PAGE 2-A



AB A review. **Oxidation** of green tea **catechins** under various conditions was chemical investigated. When a mixture of (-)-**epicatechin** (EC) and (-)-epigallocatechin (EGCG) was treated with homogenate of fresh tea leaves, theasinensin A and a new dimeric product 1, derived by Diels-Alder coupling of EGCG with its **oxidation** product, were obtained. On the other hand, auto-**oxidation** of theaflavin (TF) at pH 7.3 gave theanaphthoquinone (TNQ) and a new flavan-3-ol tetramer (2) having a bicyclooctane skeleton, which was also a Diels-Alder product of dehydro-TF and dihydro-TNQ. The results suggested that formation of quinehydrone type n-n complexes is important in the **catechin oxidation**. In the **process** of TF production from EC and (-)-epigallocatechin (EGC) by banana polyphenoloxidase, EC was rapidly **oxidized** to EC-quinone and the resulting EC-quinone **oxidized** EGC to EGC-quinone. After EGC was completely consumed, the EC-quinone began to **oxidize** TF to TNQ. This coupled **oxidation** mechanism was shown by rapid decrease of EGC in the presence of EC and the sparing of EC at the beginning of the reaction. In addition, when the reaction was conducted in the presence of glutathione, about 80% of EGC was not **oxidized**, even though EC was completely converted to its glutathione conjugates. Furthermore, it was also suggested that theasinensins (TS), **dimers** of EGC or EGCG found in black tea, were generated by reduction of corresponding quinones

(TS-quinones) formed by stereospecific coupling of two EGC-quinones. The presence of TS-quinones in the homogenate of fresh tea leaves was also confirmed by isolation of their phenazine derivs. These results suggested that thearubigins, the uncharacterized heterogeneous polymer of black tea, are probably produced by polymerization the quinones derived from **catechins** and their metabolites during tea fermentation

L10 ANSWER 7 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2002:831351 CAPLUS

DOCUMENT NUMBER: 138:72278

TITLE: Structures of two new **oxidation** products of green tea polyphenols generated by model tea fermentation

AUTHOR(S): Tanaka, Takashi; Mine, Chie; Kouno, Isao

CORPORATE SOURCE: Graduate School of Biomedical Sciences, Department of Molecular Medicinal Sciences, Nagasaki University, Nagasaki, 852-8521, Japan

SOURCE: Tetrahedron (2002), 58(43), 8851-8856

CODEN: TETRAB; ISSN: 0040-4020

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 490-46-0, (-)-**Epicatechin** 970-74-1,

(-)-Epigallocatechin 989-51-5, (-)-Epigallocatechin 3-O-gallate


RL: BSU (Biological study, unclassified); BIOL (Biological study)

(structures of two new **oxidation** products of green tea polyphenols generated by model tea fermentation)

RN 490-46-0 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-, (2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




RN 970-74-1 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 3,4-dihydro-2-(3,4,5-trihydroxyphenyl)-, (2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).


10/783,801



RN 989-51-5 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, (2R,3R)-3,4-dihydro-5,7-dihydroxy-2-(3,4,5-trihydroxyphenyl)-2H-1-benzopyran-3-yl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



IT 4670-05-7, Theaflavin

RL: FMU (Formation, unclassified); FORM (Formation, nonpreparative)


(structures of two new **oxidation** products of green tea

polyphenols generated by model tea fermentation)

RN 4670-05-7 CAPLUS

CN 5H-Benzocyclohepten-5-one, 1,8-bis[(2R,3R)-3,4-dihydro-3,5,7-trihydroxy-2H-1-benzopyran-2-yl]-3,4,6-trihydroxy- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



AB To clarify the **oxidation** mechanism of green tea **catechins** during tea fermentation, pure **catechins** were **oxidized** with a **catechin**-free homogenate of tea leaf. **Oxidation** of a mixture of (-)-**epicatechin** and (-)-epigallocatechin yielded a new metabolite, named dehydrotheaflavin, produced by the **oxidation** of a benzotropolone moiety of the black tea pigment theaflavin. Similar **oxidation** of a mixture of (-)-**epicatechin** and (-)-epigallocatechin 3-O-gallate afforded a new **dimer** of (-)-epigallocatechin 3-O-gallate, which was generated by the **oxidn** . and cycloaddn. of two pyrogallol rings. Structures were determined by spectroscopic **method**, and the **oxidation** mechanisms for the formation of the products were proposed.

REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 8 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2002:185102 CAPLUS

DOCUMENT NUMBER: 136:247439

TITLE: **Process for preparing**
4 α -aryl substituted **epicatechin**
derivatives

INVENTOR(S): Kozikowski, Alan P.; Romanczyk, Leo J., Jr.;
Tueckmantel, Werner

PATENT ASSIGNEE(S): Mars, Inc., USA

SOURCE: PCT Int. Appl., 31 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002020506	A2	20020314	WO 2001-US26175	20010821
WO 2002020506	A3	20030206		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF,			

10/783,801

BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG
US 6476241 B1 20021105 US 2000-655360 20000905
CA 2421513 AA 20020314 CA 2001-2421513 20010821
AU 2001083472 A5 20020322 AU 2001-83472 20010821
EP 1317437 A2 20030611 EP 2001-962277 20010821
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR
JP 2004508362 T2 20040318 JP 2002-525127 20010821
US 2003100775 A1 20030529 US 2002-214830 20020808
US 6720432 B2 20040413
US 2005014958 A1 20050120 US 2004-783801 20040220
PRIORITY APPLN. INFO.: US 2000-655360 A 20000905
WO 2001-US26175 W 20010821
US 2002-214830 A3 20020808

OTHER SOURCE(S): CASREACT 136:247439; MARPAT 136:247439


IT 89385-19-3P 223387-39-1P 299412-40-1P
301539-02-6P 330936-12-4P 330936-13-5P
330936-15-7P 330936-17-9P 330936-23-7P
330936-24-8P 330936-25-9P 330936-26-0P
330936-27-1P 330936-28-2P 330936-30-6P
330936-31-7P 403651-95-6P 403651-96-7P
403820-84-8P

RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(methods for the preparation of 4 α -aryl substituted **epicatechin** derivs.)

RN 89385-19-3 CAPLUS

CN 2H-1-Benzopyran, 2-[3,4-bis(phenylmethoxy)phenyl]-8-bromo-3,4-dihydro-3,5,7-tris(phenylmethoxy)-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




RN 223387-39-1 CAPLUS

CN 2H-1-Benzopyran, 2-[3,4-bis(phenylmethoxy)phenyl]-8-bromo-3,4-dihydro-3,5,7-tris(phenylmethoxy)-, (2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).


10/783,801



RN 299412-40-1 CAPLUS

CN Silane, [[(2R,3R)-2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-5,7-bis(phenylmethoxy)-2H-1-benzopyran-3-yl]oxy] (1,1-dimethylethyl)dimethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 301539-02-6 CAPLUS

CN 2H-1-Benzopyran, 2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-3,5,7-tris(phenylmethoxy)-, (2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).


10/783,801



RN 330936-12-4 CAPLUS

CN 2H-1-Benzopyran-4-ol, 2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-3,5,7-tris(phenylmethoxy)-, (2R,3R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 330936-13-5 CAPLUS

CN 4H-1-Benzopyran-4-one, 2-[3,4-bis(phenylmethoxy)phenyl]-2,3-dihydro-3,5,7-tris(phenylmethoxy)-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




10/783,801

RN 330936-15-7 CAPLUS

CN 2H-1-Benzopyran, 2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-3,5,7-tris(phenylmethoxy)-4-(2,4,6-trimethoxyphenyl)-, (2R,3R,4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 330936-17-9 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-[3,4-bis(acetyloxy)phenyl]-3,4-dihydro-4-(2,4,6-trimethoxyphenyl)-, triacetate, (2R,3R,4R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




RN 330936-23-7 CAPLUS

10/783,801


CN 4,8'-Bi-2H-1-benzopyran, 2,2'-bis[3,4-bis(phenylmethoxy)phenyl]-3,3',4,4'-tetrahydro-3,3',5,5',7,7'-hexakis(phenylmethoxy)-, (2R,2'R,3R,3'R,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

PAGE 1-A





PAGE 2-A



RN 330936-24-8 CAPLUS



CN Benzoic acid, 3,4,5-tris(phenylmethoxy)-, (2R,2'R,3R,3'R,4S)-2,2'-bis[3,4-bis(phenylmethoxy)phenyl]-3,3',4,4'-tetrahydro-3-hydroxy-5,5',7,7'-tetrakis(phenylmethoxy)[4,8'-bi-2H-1-benzopyran]-3'-yl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



RN 330936-25-9 CAPLUS
 CN Benzoic acid, 3,4,5-trihydroxy-, (2R,2'R,3R,3'R,4S)-2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-3,5,5',7,7'-pentahydroxy[4,8'-bi-2H-1-benzopyran]-3'-yl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 330936-26-0 CAPLUS
 CN 2H-1-Benzopyran-4-ol, 2-[3,4-bis(phenylmethoxy)phenyl]-3-[[[1,1-dimethylethyl)dimethylsilyl]oxy]-3,4-dihydro-5,7-bis(phenylmethoxy)-, (2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


10/783,801



RN 330936-27-1 CAPLUS

CN 4H-1-Benzopyran-4-one, 2-[3,4-bis(phenylmethoxy)phenyl]-3-[[(1,1-dimethylethyl)dimethylsilyl]oxy]-2,3-dihydro-5,7-bis(phenylmethoxy)-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




RN 330936-28-2 CAPLUS

CN Silane, [[(2R,3R)-2-[3,4-bis(phenylmethoxy)phenyl]-8-bromo-3,4-dihydro-5,7-bis(phenylmethoxy)-2H-1-benzopyran-3-yl]oxy] (1,1-dimethylethyl)dimethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

10/783,801





RN 330936-30-6 CAPLUS

CN Silane, [[(2R,2'R,3R,3'R,4S)-2,2'-bis[3,4-bis(phenylmethoxy)phenyl]-3,3',4,4'-tetrahydro-5,7-bis(phenylmethoxy)[4,8'-bi-2H-1-benzopyran]-3,3'-diyl]bis(oxy)]bis[(1,1-dimethylethyl)dimethyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).


PAGE 1-A





RN 330936-31-7 CAPLUS
 CN [4,8'-Bi-2H-1-benzopyran]-3,3'-diol, 2,2'-bis[3,4-bis(phenylmethoxy)phenyl]-3,3',4,4'-tetrahydro-5,5',7,7'-tetrakis(phenylmethoxy)-, (2R,2'R,3R,3'R,4S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 403651-95-6 CAPLUS
 CN 2H-1-Benzopyran-4-ol, 2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-3,5,7-tris(phenylmethoxy)-4-(2,4,6-trimethoxyphenyl)-, (2R,3S,4R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).


10/783,801



RN 403651-96-7 CAPLUS
 CN [4,8'-Bi-2H-1-benzopyran]-4-ol, 2,2'-bis[3,4-bis(phenylmethoxy)phenyl]-
 3,3'-bis[[(1,1-dimethylethyl)dimethylsilyl]oxy]-3,3',4,4'-tetrahydro-
 5,5',7,7'-tetrakis(phenylmethoxy)-, (2R,2'R,3S,3'R,4S)- (9CI) (CA INDEX
 NAME)


Absolute stereochemistry. Rotation (-).

PAGE 1-A



10/783,801


PAGE 2-A




RN 403820-84-8 CAPLUS
CN [4,8'-Bi-4H-1-benzopyran]-4-ol, 2,2'-bis[3,4-bis(phenylmethoxy)phenyl]-
2,2',3,3'-tetrahydro-3,3',5,5',7,7'-hexakis(phenylmethoxy)-,
(2R,2'R,3S,3'R,4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

PAGE 1-A



PAGE 2-A



IT 67253-04-7P 67253-05-8P
RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP
(Preparation)


10/783,801

(methods for the preparation of 4 α -aryl
substituted **epicatechin** derivs..)

RN 67253-04-7 CAPLUS

CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-
dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3R,3'R,4S)- (9CI) (CA
INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 67253-05-8 CAPLUS

CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis[3,4-
bis(acetyloxy)phenyl]-3,3',4,4'-tetrahydro-, hexaacetate,
(2R,2'R,3R,3'R,4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



10/783,801

IT 87292-49-7 223387-36-8 256236-21-2


RL: RCT (Reactant); RACT (Reactant or reagent)

(methods for the preparation of 4 α -aryl
substituted epicatechin derivs.)

RN 87292-49-7 CAPLUS

CN 2H-1-Benzopyran-3-ol, 2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-5,7-
bis(phenylmethoxy)-, (2R,3R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 223387-36-8 CAPLUS

CN 2H-1-Benzopyran-3-ol, 2-[3,4-bis(phenylmethoxy)phenyl]-8-bromo-3,4-dihydro-
5,7-bis(phenylmethoxy)-, (2R,3R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).




RN 256236-21-2 CAPLUS

CN 2H-1-Benzopyran-3-ol, 2-[3,4-bis(phenylmethoxy)phenyl]-8-bromo-3,4-dihydro-
5,7-bis(phenylmethoxy)-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



GI



I


AB **Process** for **preparing** a 4 α -aryl substituted **epicatechin** derivative including 4 α ,8- **epicatechin dimers** such as I (R₁,R₃ = H, acetyl, protected galloyl, galloyl; R₂ = H, benzyl, acetyl), is disclosed which comprises the steps of: (a) protecting C-3 hydroxyl group of 5,7,3',4'-tetra-O-benzylepicatechin; (b) **oxidizing** the 4-position of the compound of step (a) to produce protected flavan-4-one; (c) reacting the compound of step (b) with aryllithium reagents, derived by halogen/metal exchange from the aryl bromides, to form C-3 protected 4-hydroxy-4-aryl **epicatechin** derivative; (d) deoxygenating the C-4 position of the compound of step (c) with tri-n-butyltin hydride and trifluoroacetic acid, to afford C-3 protected 4 α -aryl-5,7,3',4'-tetra-O-benzylepicatechin. Thus, **epicatechin-4 α ,8-(3-O-galloylepicatechin)** I (R₁, R₂ = H; R₃ = galloyl) was **prepared** in a multistep synthetic sequence starting from 5,7,3',4'-tetra-O-benzylepicatechin, 5,7,3',4'-tetra-O-benzyl-8-bromoepicatechin, and tri-O-benzyl gallic acid.

10/783,801

ACCESSION NUMBER: 2001:408634 CAPLUS
DOCUMENT NUMBER: 135:45464
TITLE: Changes of (+)-**catechin** antioxidant activity during enzymic **oxidation**
AUTHOR(S): Sosnowska, Dorota
CORPORATE SOURCE: Inst. Biochem. Tech., Politech. Lodzka, Lodz, 90-924, Pol.
SOURCE: Zywnosc (2000), 7(3, Supl.), 69-76
CODEN: ZYWNFL
PUBLISHER: Polskie Towarzystwo Technologow Zywnosci, Oddzial Malopolski
DOCUMENT TYPE: Journal
LANGUAGE: Polish

IT 154-23-4, +**Catechin**
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(antioxidant activity of (+)-**catechin** changes during enzymic **oxidation** with apple polyphenol oxidase and tyrosinase)
RN 154-23-4 CAPLUS
CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




AB The effects of enzymic **oxidation** of com. pure (+)-**catechin** on its antioxidant activity were investigated. The **oxidation** of (+)-**catechin**, a common component of fruits and vegetables, with apple polyphenol oxidase (PPO) and tyrosinase was studied at 25°C in solns. at pH 4 and 7. The antioxidant activity was determined by 2,2'-azinobis-3(ethylbenzothiazoline-6-sulfonic acid) (ABTS) and 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging. The (+)-**catechin** and its reaction products were monitored by HPLC and TLC. The rate of **oxidation** reaction in the presence of PPO at pH 7 was higher than at pH 4. The reaction products of **catechin oxidation** were **dimers** and tetramers. The **oxidation** of **catechin** with tyrosinase was slower than with PPO. The antioxidant activity of **catechin** after 2 h **oxidation** (PPO at pH 7) was decreased .apprx.25% as determined by the DPPH **method** (expressed as IC50) and .apprx.12% as determined by the ABTS **method** (expressed as TEAC - Trolox Equivalent antioxidant capacity) despite large decrease in the content of monomeric **catechin** (57%). This indicates that (+)-**catechin** enzymic **oxidation** produces compds. with antioxidant activities.

L10 ANSWER 10 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 2000:830787 CAPLUS

10/783,801

DOCUMENT NUMBER: 134:125908
TITLE: Evaluation of the Antioxidant and Pro-oxidant Effects of Tea **Catechin** Oxypolymers
AUTHOR(S): Li, Chunmei; Xie, Bijun
CORPORATE SOURCE: Department of Food Science and Technology, Huazhong Agriculture University, Wuhan, Hubei, 430070, Peop. Rep. China
SOURCE: Journal of Agricultural and Food Chemistry (2000), 48(12), 6362-6366
CODEN: JAFCAU; ISSN: 0021-8561
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
IT **989-51-5D**, Epigallocatechin gallate, oxypolymers
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); BIOL (Biological study); RACT (Reactant or reagent)
(antioxidant and pro-oxidant effects of tea **catechin** oxypolymers)
RN 989-51-5 CAPLUS
CN Benzoic acid, 3,4,5-trihydroxy-, (2R,3R)-3,4-dihydro-5,7-dihydroxy-2-(3,4,5-trihydroxyphenyl)-2H-1-benzopyran-3-yl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



AB Tea **catechin** oxypolymers (TCOP) were **prepared** by **oxidizing** tea **catechin** (TC, the content of epigallocatechin gallate (EGCG) was >85%) with H₂O₂. TCOP was a mixture of polymers mainly **dimers** and trimers of EGCG. Their antioxidant and pro-oxidant effects were tested using a deoxyribose assay, a photoredn. of NBT assay, a lipoxxygenase assay, a POV assay, and animal tests. The scavenging effects of TCOP to both the hydroxyl radical and superoxide radical were stronger than that of TC, and also they had no pro-oxidant effect; the rate constant for reactions of TC and TCOP for hydroxyl radical were 1.0×10^{10} and $(1.4-2.8) \times 10^{10} \text{ M}^{-1} \text{ S}^{-1}$, resp. TCOP can inhibit lipid peroxidn. and lipoxxygenase effectively, and it also can activate red cell SOD and reduce the MDA content in serum of mice very significantly. These results suggested that the antioxidant activity of TCOP was not less than or even more notable than that of TC.
REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 11 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2000:553825 CAPLUS

DOCUMENT NUMBER: 133:335114

TITLE: Antioxidant properties of anthocyanins and tannins: a mechanistic investigation with **catechin** and the 3',4',7-trihydroxyflavylium ion

AUTHOR(S): Dangles, Olivier; Fargeix, Guillaume; Dufour, Claire

CORPORATE SOURCE: UMR-CNRS 5078, Bat. 303, Universite Claude

Bernard-Lyon I, Villeurbanne, 69622, Fr.

SOURCE: Perkin 2 (2000), (8), 1653-1663

CODEN: PRKTFO

PUBLISHER: Royal Society of Chemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 133:335114

IT 154-23-4, **Catechin**


RL: PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); PROC (Process); RACT (Reactant or reagent)

(antioxidant properties of anthocyanins and tannins: a mechanistic investigation with **catechin** and 3',4',7-trihydroxyflavylium ion)


RN 154-23-4 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



GI



AB Plant polyphenols act as antioxidants mainly by trapping reactive oxygen species and by regenerating endogenous membrane-bound α -tocopherol (vitamin E). In both **processes** polyphenols are **oxidized**. Hence, knowledge of the **oxidation** mechanisms of polyphenols is important for an understanding of their antioxidant activity at the mol. level. This work focuses on anthocyanins (pigments) and flavanols (tannins), two important classes of polyphenols which are both relatively abundant in human diet. The **oxidation** of the 3',4',7-trihydroxyflavylium ion (I) and **catechin** (II), resp. taken as models for anthocyanins and tannins, has been investigated. From kinetic data and partial product anal., the mechanisms for the reactions of I and II with sodium periodate and DPPH, a H atom-abstracting radical, are proposed. Both polyphenols are shown to form o-quinone intermediates upon H atom abstraction and subsequent radical disproportionation. In the case of II, the quinone and a second mol. of antioxidant quickly couple to form **dimers**. By contrast, I is extensively degraded into coumarins by repeating sequences of **oxidation**-solvent addition, which consume several equivalent of oxidants. In aqueous solns., I is typically a mixture of colored and colorless forms. The latter (chalcones) are also shown to take part in the antioxidant activity.

REFERENCE COUNT: 73 THERE ARE 73 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 12 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2000:539327 CAPLUS

DOCUMENT NUMBER: 133:222006

TITLE: **Epicatechin** in human plasma: in vivo determination and effect of chocolate consumption on plasma **oxidation** status

AUTHOR(S): Rein, Dietrich; Lotito, Silvina; Holt, Roberta R.; Keen, Carl L.; Schmitz, Harold H.; Fraga, Cesar G.

CORPORATE SOURCE: Department of Nutrition, University of California, Davis, CA, 95616, USA

SOURCE: Journal of Nutrition (2000), 130(8S), 2109S-2114S
CODEN: JONUAI; ISSN: 0022-3166

PUBLISHER: American Society for Nutritional Sciences

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 154-23-4, **Catechin**


RL: ANT (Analyte); ANST (Analytical study)

(dietary chocolate effects on **epicatechin** blood plasma kinetics and **oxidation** status in humans)

RN 154-23-4 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).

IT **490-46-0, Epicatechin**


RL: ANT (Analyte); BPR (Biological process); BSU (Biological study, unclassified); FFD (Food or feed use); ANST (Analytical study); BIOL (Biological study); PROC (Process); USES (Uses)

(dietary chocolate effects on **epicatechin** blood plasma kinetics and **oxidation** status in humans)

RN 490-46-0 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-, (2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



AB Diets rich in plant foods have been associated with decreased risk for specific disease **processes** and certain chronic diseases. In addition to essential macronutrients and micronutrients, flavonoids in plant foods may have health-enhancing properties. Chocolate is rich in the flavan-3-ol **epicatechin** and procyanidin oligomers. The bioavailability and biol. effects of the chocolate flavonoids are poorly understood. HPLC coupled with electrochem. (coulometric) detection was used to determine the physiolo. levels of **epicatechin**, **catechin**, and **epicatechin dimers** in human blood plasma. This **method** allows to determine levels as low as 20 pg (69 fmol) of **epicatechin** (1 nmol/L blood plasma). The absorption of **epicatechin** from an 80-g dose of semisweet procyanidin-rich chocolate was evaluated using this **method** in 13 healthy adult humans. By 2 h after ingestion, there was a 12-fold increase in plasma **epicatechin** levels from 22 to 257 nmol/L. Consistent with the antioxidant properties of **epicatechin**, within the same 2-h period there was a 31% increase in plasma total antioxidant capacity and 40% decrease in the plasma 2-thiobarbituric acid-reactive substance levels. Plasma **epicatechin** and plasma

antioxidant capacity approached baseline values by 6 h after chocolate ingestion. Thus, it is possible to determine basal levels of **epicatechin** in blood plasma. The consumption of chocolate can increase blood plasma **epicatechin** and decrease plasma baseline **oxidation** product concns.

REFERENCE COUNT: 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 13 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2000:128621 CAPLUS

DOCUMENT NUMBER: 132:278340

TITLE: Xanthylum salts formation involved in wine color changes

AUTHOR(S): Es-Safi, Nour-Eddine; Le Guerneve, Christine; Fulcrand, Helene; Cheynier, Veronique; Moutounet, Michel

CORPORATE SOURCE: Vigne, Unite de Recherche Biopolymeres et Aromes, ISVV-INRA Institut des Produits, Montpellier, 34060, Fr.

SOURCE: International Journal of Food Science and Technology (2000), 35(1), 63-74

CODEN: IJFTEZ; ISSN: 0950-5423

PUBLISHER: Blackwell Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 246181-58-8


RL: FMU (Formation, unclassified); FORM (Formation, nonpreparative)
(xanthylum salts formation involved in wine color changes)


RN 246181-58-8 CAPLUS

CN 2H-1-Benzopyran-8-acetic acid, 2-(3,4-dihydroxyphenyl)- α -[(2R,3S)-2-(3,4-dihydroxyphenyl)-3,4-dihydro-3,5,7-trihydroxy-2H-1-benzopyran-8-yl]-3,4-dihydro-3,5,7-trihydroxy-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A





IT 253120-56-8 253120-57-9 253120-58-0


RL: FMU (Formation, unclassified); RCT (Reactant); FORM (Formation, nonpreparative); RACT (Reactant or reagent)

(xanthylum salts formation involved in wine color changes)

RN 253120-56-8 CAPLUS

CN 2H-1-Benzopyran-6-acetic acid, 2-(3,4-dihydroxyphenyl)- α -[(2R, 3S)-2-(3,4-dihydroxyphenyl)-3,4-dihydro-3,5,7-trihydroxy-2H-1-benzopyran-8-yl]-3,4-dihydro-3,5,7-trihydroxy-, (α S, 2R, 3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 253120-57-9 CAPLUS

CN 2H-1-Benzopyran-6-acetic acid, 2-(3,4-dihydroxyphenyl)- α -[(2R, 3S)-2-(3,4-dihydroxyphenyl)-3,4-dihydro-3,5,7-trihydroxy-2H-1-benzopyran-8-yl]-3,4-dihydro-3,5,7-trihydroxy-, (α R, 2R, 3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


10/783,801



RN 253120-58-0 CAPLUS

CN 2H-1-Benzopyran-6-acetic acid, 2-(3,4-dihydroxyphenyl)- α -[(2R,3S)-2-(3,4-dihydroxyphenyl)-3,4-dihydro-3,5,7-trihydroxy-2H-1-benzopyran-6-yl]-3,4-dihydro-3,5,7-trihydroxy-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 154-23-4, (+)-**Catechin**

RL: RCT (Reactant); RACT (Reactant or reagent)
(xanthylum salts formation involved in wine color changes)

RN 154-23-4 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



AB The reaction of (+)-**catechin** in wine-like model solution was investigated. First appearance of colorless dimeric compds. consisting of 2 flavanol units linked by carboxymethine bridge was observed. Their isolation and further incubation was found to yield 2 types of yellowish pigments showing visible absorption maxima at 440 and 460 nm, resp. Mass spectroscopy (MS) spectral anal. showed that the 1st type were xanthylum salt pigments formed by dehydration of the colorless compds. followed by an **oxidation process**. The loss of a water mol. was shown to take place between 2 A ring hydroxyl groups of the colorless **dimers**. The 2nd type were shown to be ester derivs. of the 1st ones. Thus, the Et ester of xanthylum salt was obtained and fully characterized by mass and NMR spectroscopy. Esterification was found to involve the colorless compound before dehydration and thus a general scheme for xanthylum salt formation was postulated. The proposed scheme constitutes a new xanthylum formation pathway as up to now only anthocyanin-flavanol reactions were supposed to form xanthylum salt derivs. during wine ageing. This work also provides new support to the contribution of xanthylum salt in color evolution observed during wine ageing which is generally expressed in an increase of absorption in the 400-500 nm, region of xanthylum salt absorption maxima.

REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 14 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 2000:113981 CAPLUS

DOCUMENT NUMBER: 132:251007

TITLE: Tyrosinase catalyzed biphenyl construction from flavan-3-ol substrates

AUTHOR(S): Van Rensburg, Werner Janse; Ferreira, Daneel; Malan, Elfranco; Steenkamp, Jacobus A.

CORPORATE SOURCE: Department of Chemistry, University of the Orange Free State, Bloemfontein, 9300, S. Afr.

SOURCE: Phytochemistry (2000), 53(2), 285-292

CODEN: PYTCAS; ISSN: 0031-9422

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 132:251007

IT 109671-61-6P 109718-27-6P

RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)


(tyrosinase catalyzed biphenyl construction from flavan-3-ol substrates)

RN 109671-61-6 CAPLUS

CN [5,8'-Bi-2H-1-benzopyran]-3,3',5',7,7',8-hexol, 2,2'-bis(3,4-


10/783,801

dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3S,3'S,5S)- (9CI) (CA INDEX NAME)



RN 109718-27-6 CAPLUS

CN [5,8'-Bi-2H-1-benzopyran]-3,3',5',7,7',8-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3S,3'S,5R)- (9CI) (CA INDEX NAME)



IT 30825-50-4P


RL: BPN (Biosynthetic preparation); RCT (Reactant); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent)
(tyrosinase catalyzed biphenyl construction from flavan-3-ol substrates)

RN 30825-50-4 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 3,4-dihydro-2-(2',4,4',5,6'-pentahydroxy[1,1'-biphenyl]-2-yl)-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

10/783,801




IT **263026-49-9P**

RL: BPN (Biosynthetic preparation); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(tyrosinase catalyzed biphenyl construction from flavan-3-ol substrates)

RN 263026-49-9 CAPLUS

CN 2H-1-Benzopyran-3-ol, 3,4-dihydro-5,7-dimethoxy-2-(2',4,4',5,6'-pentamethoxy[1,1'-biphenyl]-2-yl)-, acetate, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT **154-23-4, Catechin**

RL: RCT (Reactant); RACT (Reactant or reagent)
(tyrosinase catalyzed biphenyl construction from flavan-3-ol substrates)

RN 154-23-4 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



AB Mushroom tyrosinase catalyzed **oxidation** of three flavan-3-ols, viz. **catechin**, fisetinidol and mesquitol, was conducted to construct biphenyl bonds. Exposure of the flavan-3-ols to tyrosinase and subsequent trapping of the o-quinone intermediates resulted in the formation of novel flavan-3-ol derivs., the structures of which were elucidated by mono- and two-dimensional 1H-NMR expts. Application of the methodol. resulted in the improved **synthesis** of the natural flavan-3-ol **dimer**, mesquitol-[5 → 8]- **catechin**, previously isolated from *Prosopis glandulosa*.

REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 15 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1999:771159 CAPLUS

DOCUMENT NUMBER: 132:122417

TITLE: Studies in Polyphenol Chemistry and Bioactivity. 1. **Preparation** of Building Blocks from (+)-**Catechin**. Procyanidin Formation.

Synthesis of the Cancer Cell Growth Inhibitor, 3-O-Galloyl-(2R,3R)-**epicatechin**-4 β ,8-[3-O-galloyl-(2R,3R)-**epicatechin**]

AUTHOR(S): Tueckmantel, Werner; Kozikowski, Alan P.; Romanczyk, Leo J., Jr.

CORPORATE SOURCE: Georgetown University Medical Center Institute for Cognitive and Computational Sciences Drug Discovery Program, Washington, DC, 20007, USA

SOURCE: Journal of the American Chemical Society (1999), 121(51), 12073-12081

CODEN: JACSAT; ISSN: 0002-7863

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 132:122417

IT 256236-26-7P 256236-27-8P 256236-28-9P 256236-29-0P


RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation) (**synthesis** of [(2R,3R)-3-O-galloylepicatechin]-4 β ,8-[(2R,3R)-3-O-galloylepicatechin] from (+)-**catechin**)

RN 256236-26-7 CAPLUS

CN Benzeneacetic acid, α -methoxy- α -(trifluoromethyl)-, (2R,3S)-2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-5,7-bis(phenylmethoxy)-2H-1-benzopyran-3-yl ester, (α S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/783,801



RN 256236-27-8 CAPLUS
CN Benzeneacetic acid, α -methoxy- α -(trifluoromethyl)-,
(2R,3S)-2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-5,7-
bis(phenylmethoxy)-2H-1-benzopyran-3-yl ester, (α R)- (9CI) (CA
INDEX NAME)


Absolute stereochemistry.



RN 256236-28-9 CAPLUS
CN Benzeneacetic acid, α -methoxy- α -(trifluoromethyl)-,
(2R,3R)-2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-5,7-
bis(phenylmethoxy)-2H-1-benzopyran-3-yl ester, (α S)- (9CI) (CA
INDEX NAME)


Absolute stereochemistry.

10/783,801




RN 256236-29-0 CAPLUS
CN Benzeneacetic acid, α -methoxy- α -(trifluoromethyl)-,
(2R,3R)-2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-5,7-
bis(phenylmethoxy)-2H-1-benzopyran-3-yl ester, (α R)- (9CI) (CA
INDEX NAME)

Absolute stereochemistry.




IT **154-23-4, (+)-Catechin**
RL: RCT (Reactant); RACT (Reactant or reagent)
(**synthesis** of [(2R,3R)-3-O-galloylepicatechin]-4 β ,8-[-
(2R,3R)-3-O-galloylepicatechin] from (+)-**catechin**)
RN 154-23-4 CAPLUS
CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-,
(2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



IT 20728-73-8P, 5,7,3',4'-Tetra-O-benzylcatechin 29106-49-8P
 , (+)-Procyanidin B2 79907-44-1P, (2R,3R)-3-O-Galloylepicatechin-
 4 β ,8-[(2R,3R)-3-O-galloylepicatechin] 87292-49-7P,
 (-)-5,7,3',4'-Tetra-O-benzylepicatechin 223387-28-8P
 223387-33-5P 256236-21-2P, (-)-5,7,3',4'-Tetra-O-benzyl-
 8-bromocatechin 256236-25-6P 256236-30-3P,
 (-)-5,7,3',4'-Tetra-O-benzyl-3-O-(3,4,5-tri-O-benzylgalloyl)
epicatechin
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (synthesis of [(2R,3R)-3-O-galloylepicatechin]-4 β ,8-[-
 (2R,3R)-3-O-galloylepicatechin] from (+)-catechin)
 RN 20728-73-8 CAPLUS
 CN 2H-1-Benzopyran-3-ol, 2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-5,7-
 bis(phenylmethoxy)-, (2R,3S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



RN 29106-49-8 CAPLUS
 CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-
 dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3R,3'R,4R)- (9CI) (CA
 INDEX NAME)

Absolute stereochemistry.

10/783,801





RN 79907-44-1 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, (2R,2'R,3R,3'R,4R)-2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-5,5',7,7'-tetrahydroxy[4,8'-bi-2H-1-benzopyran]-3,3'-diyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

PAGE 1-A






RN 87292-49-7 CAPLUS

CN 2H-1-Benzopyran-3-ol, 2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-5,7-bis(phenylmethoxy)-, (2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




RN 223387-28-8 CAPLUS

CN [4,8'-Bi-2H-1-benzopyran]-3,3'-diol, 2,2'-bis[3,4-bis(phenylmethoxy)phenyl]-3,3',4,4'-tetrahydro-5,5',7,7'-tetrakis(phenylmethoxy)-, (2R,2'R,3R,3'R,4R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).

10/783,801





RN 223387-33-5 CAPLUS

CN Benzoic acid, 3,4,5-tris(phenylmethoxy)-, (2R,2'R,3R,3'R,4R)-2,2'-bis[3,4-bis(phenylmethoxy)phenyl]-3,3',4,4'-tetrahydro-5,5',7,7'-tetrakis(phenylmethoxy)[4,8'-bi-2H-1-benzopyran]-3,3'-diyl ester (9CI)
(CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

PAGE 1-A






RN 256236-21-2 CAPLUS

CN 2H-1-Benzopyran-3-ol, 2-[3,4-bis(phenylmethoxy)phenyl]-8-bromo-3,4-dihydro-5,7-bis(phenylmethoxy)-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




RN 256236-25-6 CAPLUS

CN 2H-1-Benzopyran-3-ol, 2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-4-(2-hydroxyethoxy)-5,7-bis(phenylmethoxy)-, (2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


10/783,801



RN 256236-30-3 CAPLUS

CN Benzoic acid, 3,4,5-tris(phenylmethoxy)-, (2R,3R)-2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-5,7-bis(phenylmethoxy)-2H-1-benzopyran-3-yl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



IT 1257-08-5P, 3-O-Galloylepicatechin 21179-21-5P,
(+)-Procyanidin B2 decaacetate 223387-36-8P,
(-)-8-Bromo-5,7,3',4'-tetra-O-benzylepicatechin 256236-23-4P
256236-24-5P 256236-31-4P


RL: SPN (Synthetic preparation); PREP (Preparation)
(**synthesis** of [(2R,3R)-3-O-galloylepicatechin]-4 β ,8-[-
(2R,3R)-3-O-galloylepicatechin] from (+)-catechin)

RN 1257-08-5 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, (2R,3R)-2-(3,4-dihydroxyphenyl)-3,4-dihydro-5,7-dihydroxy-2H-1-benzopyran-3-yl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).

10/783,801



RN 21179-21-5 CAPLUS
CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis[3,4-bis(acetyloxy)phenyl]-3,3',4,4'-tetrahydro-, hexaacetate, (2R,2'R,3R,3'R,4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



RN 223387-36-8 CAPLUS
CN 2H-1-Benzopyran-3-ol, 2-[3,4-bis(phenylmethoxy)phenyl]-8-bromo-3,4-dihydro-5,7-bis(phenylmethoxy)-, (2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).


10/783,801



RN 256236-23-4 CAPLUS

CN 2H-1-Benzopyran-3-ol, 2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-4-methoxy-5,7-bis(phenylmethoxy)-, (2R,3R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 256236-24-5 CAPLUS

CN 2H-1-Benzopyran-3-ol, 2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-2,4-dimethoxy-5,7-bis(phenylmethoxy)-, (3R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



10/783,801

RN 256236-31-4 CAPLUS
CN 2H-1-Benzopyran-3,4-diol, 2-[3,4-bis(phenylmethoxy)phenyl]-3,4-dihydro-5,7-bis(phenylmethoxy)-, (2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB A project has been initiated to **synthesize** proanthocyanidin oligomers found in cocoa. Natural, readily available (+)-**catechin** was transformed into 5,7,3',4'-tetra-O-benzyl-(-)-**epicatechin** (I) by (a) benzylation of the phenolic oxygens; (b) **oxidation** of the 3-alc. to ketone by the Dess-Martin periodinane; and (c) reduction with lithium tri-sec-butylborohydride (L-Selectride) in the presence of LiBr. The additive diminishes the extent of ketone enolization while maintaining a stereoselectivity of $\geq 200:1$. **Oxidation** of I with DDQ was performed best from the standpoint of product purification if ethylene glycol was used as the nucleophilic trapping agent. The resulting ether II was condensed with I using TiCl₄ to give a good yield of benzyl-protected **epicatechin-4 β ,8-epicatechin** (octa-O-benzylprocyanidin B2, III) as the sole dimeric product. Hydrogenolysis of III yielded procyanidin B2 in the first enantiospecific **synthesis** of this natural product which employs protected intermediates and thereby allows the necessary product separation after the condensation step to be performed on nonpolar, nonsensitive intermediates. Acylation of III with tri-O-benzylgalloyl chloride followed by hydrogenolysis gave access to the title bis-gallate IV [R = COC₆H₂(OH)_{3-3,4,5}]. This constitutes the first **synthesis** of this natural product, a compound notable for its PKC-inhibitory and anticancer activity.

REFERENCE COUNT: 201 THERE ARE 201 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L10 ANSWER 16 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1999:195757 CAPLUS

DOCUMENT NUMBER: 131:5123

TITLE: Antioxidant Chemistry of Green Tea **Catechins**
. Identification of Products of the Reaction of

AUTHOR(S): (-)-Epigallocatechin Gallate with Peroxyl Radicals
Valcic, Susanne; Muders, Annette; Jacobsen, Neil E.;
Liebler, Daniel C.; Timmermann, Barbara N.

CORPORATE SOURCE: Department of Pharmacology Toxicology College of
Pharmacy, The University of Arizona, Tucson, AZ,
85721, USA

SOURCE: Chemical Research in Toxicology (1999), 12(4), 382-386
CODEN: CRTOEC; ISSN: 0893-228X

PUBLISHER: American Chemical Society


DOCUMENT TYPE: Journal

LANGUAGE: English

IT 989-51-5, (-)-Epigallocatechin gallate
RL: PRP (Properties); RCT (Reactant); RACT (Reactant or reagent)
(antioxidant chemical of (-)-epigallocatechin gallate from green tea)

RN 989-51-5 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, (2R,3R)-3,4-dihydro-5,7-dihydroxy-2-
(3,4,5-trihydroxyphenyl)-2H-1-benzopyran-3-yl ester (9CI) (CA INDEX NAME)



AB (-)-Epigallocatechin gallate (EGCG), isolated from green tea, displays antioxidant properties and is thought to act as an antioxidant in biol. systems. However, the specific mechanisms of its antioxidant actions remain unclear. The authors have isolated and identified for the first time two reaction products of EGCG derived from its reaction with peroxy radicals generated by thermolysis of the initiator 2,2'-azobis(2,4-dimethylvaleronitrile) (AMVN). The products include a seven-membered B-ring anhydride and a novel **dimer**. The identification of these products provides the first unambiguous proof that the principal site of antioxidant reactions on the EGCG mol. is the trihydroxyphenyl B ring, rather than the 3-galloyl moiety. In contrast to phenoxyl radicals from simple phenolic antioxidants, an initially formed EGCG phenoxyl radical apparently does not form stable addition products with AMVN-derived peroxy radicals. Characteristic reaction products may provide novel markers for EGCG antioxidant reactions in living systems.

REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 17 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1999:157268 CAPLUS

10/783,801

DOCUMENT NUMBER: 130:301542
TITLE: Biochemical activities of extracts from *Hypericum perforatum*. 3rd communication. Modulation of peroxidase activity as a simple **method** for standardization
AUTHOR(S): Schempp, Harald; Denke, Andrea; Mann, Elke; Schneider, Werner; Elstner, Erich F.
CORPORATE SOURCE: Labor Angewandte Biochemie, Technische Universitaet Muenchen, Freising, D-85350, Germany
SOURCE: Arzneimittel-Forschung (1999), 49(2), 115-119
CODEN: ARZNAD; ISSN: 0004-4172
PUBLISHER: Editio Cantor Verlag
DOCUMENT TYPE: Journal
LANGUAGE: English

IT 154-23-4 490-46-0, **Epicatechin**


RL: ANT (Analyte); BOC (Biological occurrence); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study); OCCU (Occurrence)

(components of *Hypericum* exts. by HPLC separation)

RN 154-23-4 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-, (2R,3S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



RN 490-46-0 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-, (2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



AB Alc. exts. from the herb "St. John's wort" (*Hypericum perforatum*) are

widely used to counteract depressive situations, where the question on the mainly active principle is still under discussion. Thus, standardization of the drug on the basis of dry matter was chosen instead of the popular leading component, hypericin. Inhibition of myeloperoxidase-catalyzed dimerization of enkephalins by Hypericum exts. was recently reported. This **method** is based on the separation and quantification of enkephalin **dimers** by HPLC. To simplify this assay myeloperoxidase could be substituted by the cheaper horseradish peroxidase and the enkephalins by the amino acid tyrosine without loss of significance. In this communication we represent a more rapid photometric **method** based on peroxidase-catalyzed indole acetic acid **oxidation** suitable for quick, simple, and economic drug standardization.

REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 18 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1998:591261 CAPLUS

DOCUMENT NUMBER: 129:188607

TITLE: Effect of malvidin-3-monoglucoside on the browning of monomeric and dimeric flavanols

AUTHOR(S): Francia-Aricha, E. M.; Rivas-Gonzalo, J. C.; Santos-Buelga, C.

CORPORATE SOURCE: Dep. Quimica Analitica, Nutricion, Bromatologia, Fac. Farmacia, Univ. Salamanca, Salamanca, E-37007, Spain

SOURCE: Zeitschrift fuer Lebensmittel-Untersuchung und -Forschung A: Food Research and Technology (1998), 207(3), 223-228

CODEN: ZLFAFA; ISSN: 1431-4630

PUBLISHER: Springer-Verlag

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 154-23-4, (+)-Catechin 490-46-0, (-)-

Epicatechin 7228-78-6, Malvidin-3-monoglucoside

23567-23-9, Procyanidin B3 **29106-49-8**, Procyanidin B2


RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(effect of malvidin-3-monoglucoside on browning of monomeric and dimeric flavanols)

RN 154-23-4 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




RN 490-46-0 CAPLUS

10/783,801

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-,
(2R,3R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 7228-78-6 CAPLUS

CN 1-Benzopyrylium, 3-(β -D-glucopyranosyloxy)-5,7-dihydroxy-2-(4-hydroxy-3,5-dimethoxyphenyl)-, chloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 23567-23-9 CAPLUS

CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3S,3'S,4S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).

10/783,801



RN 29106-49-8 CAPLUS
 CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3R,3'R,4R,4'R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




AB The behavior of (+)-**catechin** (cat), (-)-**epicatechin** (epi) and the procyanidin **dimers** B2 [epi-(4 → 8)-epi] and B3 [cat-(4 → 8)-cat], alone and in the presence of malvidin-3-monoglucoside, was studied in hydro-alc. solns. at pH 3.2 maintained at 32° for 120 days. Browning was observed both in the presence and absence of the anthocyanin, except in the solution containing only cat. The formation of pigments showing maximum absorption in the visible region of 440-460 nm was detected. The majority of these new pigments did

not require the anthocyanin for their formation, but rather were derived directly from the flavanol. Some specific pigments showing similar spectra appeared in each of the solns. containing both a flavanol and the anthocyanin, and their production was attributed to a condensation **process** between the two latter compds. The cat units showed a greater tendency to condense with the anthocyanin, while the epi units were more sensitive to chemical **oxidation**. The formation of pigments which showed maximum absorbance in the visible region .apprx.530-545 nm, resulting from the condensation between the flavanol and the anthocyanin mediated by acetaldehyde, was also detected. As no acetaldehyde was added to the solns., its presence was attributed to the **oxidation** of ethanol coupled to autoxidn. of the catechol rings of the flavanols.

L10 ANSWER 19 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN
 ACCESSION NUMBER: 1998:66516 CAPLUS
 DOCUMENT NUMBER: 128:139871
 TITLE: Incidence of Molecular Structure in **Oxidation**
 of Grape Seeds Procyanidins
 AUTHOR(S): de Freitas, Victor A. P.; Glories, Yves; Laguerre,
 Michel
 CORPORATE SOURCE: Departamento de Quimica, Faculdade de Ciencias da
 Universidade do Porto, Oporto, 4150, Port.
 SOURCE: Journal of Agricultural and Food Chemistry (1998),
 46(2), 376-382
 CODEN: JAFCAU; ISSN: 0021-8561
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 154-23-4, (+)-**Catechin** 490-46-0, (-)-
Epicatechin 12798-58-2, Procyanidin B6
 12798-59-3, Procyanidin B7 12798-60-6, Procyanidin B8
 20315-25-7, Procyanidin B1 23567-23-9, Procyanidin B3
 29106-49-8, Procyanidin B2 29106-51-2, Procyanidin B4
 37064-30-5, Procyanidin C1 73086-04-1 89064-33-5
 97233-64-2 201412-28-4 202473-91-4
 RL: BPR (Biological process); BSU (Biological study, unclassified); PEP
 (Physical, engineering or chemical process); BIOL (Biological study); PROC
 (Process)
 (incidence of mol. structure in **oxidation** of grape seeds
 procyanidins)
 RN 154-23-4 CAPLUS
 CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-,
 (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




10/783,801

RN 490-46-0 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol; 2-(3,4-dihydroxyphenyl)-3,4-dihydro-,
(2R,3R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 12798-58-2 CAPLUS

CN [4,6'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-
dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3S,3'S,4R)- (9CI) (CA
INDEX NAME)

Absolute stereochemistry.




RN 12798-59-3 CAPLUS

CN [4,6'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-
dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3R,3'S,4S)- (9CI) (CA
INDEX NAME)


Absolute stereochemistry.

10/783,801



RN 12798-60-6 CAPLUS
CN [4,6'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3S,3'R,4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 20315-25-7 CAPLUS
CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3R,3'S,4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/783,801



RN 23567-23-9 CAPLUS
CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3S,3'S,4S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 29106-49-8 CAPLUS
CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3R,3'R,4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/783,801





RN 29106-51-2 CAPLUS
 CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3S,3'R,4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



RN 37064-30-5 CAPLUS
 CN [4,8':4',8''-Ter-2H-1-benzopyran]-3,3',3'',5,5',5'',7,7',7''-nonol, 2,2',2''-tris(3,4-dihydroxyphenyl)-3,3',3'',4,4',4''-hexahydro-, (2R,2'R,2''R,3R,3'R,3''R,4R,4'S)- (9CI) (CA INDEX NAME)



Absolute stereochemistry. Rotation (+).



RN 73086-04-1 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, (2R,2'R,3R,3'R,4R)-2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-3,5,5',7,7'-pentahydroxy[4,8'-bi-2H-1-benzopyran]-3'-yl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 89064-33-5 CAPLUS



CN Benzoic acid, 3,4,5-trihydroxy-, (2R,2'R,3S,3'R,4S)-2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-3,5,5',7,7'-pentahydroxy[4,8'-bi-2H-1-benzopyran]-3'-yl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.





RN 97233-64-2 CAPLUS
 CN [4,8':4',8''-Ter-2H-1-benzopyran]-3,3',3'',5,5',5'',7,7',7''-nonol,
 2,2',2''-tris(3,4-dihydroxyphenyl)-3,3',3'',4,4',4''-hexahydro-,
 (2R,2'R,2''R,3S,3'S,3''R,4S,4'R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.





RN 201412-28-4 CAPLUS
 CN Benzoic acid, 3,4,5-trihydroxy-, 3-[(3,4-dihydroxybenzoyl)oxy]-2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-5,5',7,7'-tetrahydroxy[4,8'-bi-2H-1-benzopyran]-3'-yl ester, [2R-[2 α ,3 α ,4 β (2'R*,3'R*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 202473-91-4 CAPLUS
 CN [4,8':4',6''-Ter-2H-1-benzopyran]-3,3',3'',5,5',5'',7,7',7''-nonol,
 2,2',2''-tris(3,4-dihydroxyphenyl)-3,3',3'',4,4',4''-hexahydro-,
 (2R,2'R,2''R,3S,3'R,3''S,4S,4'S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



AB The kinetics of decomposition of the following flavan-3-ol derivs. isolated from grape seeds under oxidative conditions by airing and using metal ion catalysis (iron and copper) are determined: (+)-**catechin** and (-)-**epicatechin**; seven natural procyanidin **dimers**, B1 [(-)-**epicatechin**-(4→8)-(+)-**catechin**], B2 [(-)-**epicatechin**-(4→8)-(-)-**epicatechin**], B3 [(+)-**catechin**-(4→8)-(+)-**catechin**], B4 [(+)-**catechin**-(4→8)-(-)-**epicatechin**], B6 [(+)-**catechin**-(4→6)-(+)-**catechin**], B7 [(-)-**epicatechin**-(4→6)-(-)-**epicatechin**], and B8 [(-)-**epicatechin**-(4→8)-(-)-**epicatechin**]; trimers, C1 [(-)-**epicatechin**-(4→8)-(-)-**epicatechin**-(4→8)-(-)-**epicatechin**], (+)-**catechin**-(4→8)-(+)-**catechin**-(4→8)-(-)-**epicatechin**, and (+)-**catechin**-(4→8)-(-)-**epicatechin**

-(4→6)-(-)- **catechin**, monogallate esters of B2 and B4 and digallate of B2, which were isolated from grape seeds. Kinetic decomposition comparisons were monitored by HPLC. The following order was found for oxidative decomposition for procyanidin **dimers**: B3 ≈ B4 > B5 ≈ B6 > B1 ≈ B2 > B8. In the conditions of this study, the gallate ester of (-)-**epicatechin** is more unstable than (-)-**epicatechin**; inversely, kinetic decompns. of dimeric procyanidins B2 and B4 are much more important than those of their gallate esters. Mol. mechanics (MM2*) and 1H NMR studies of dimeric 3-O-gallate structures show a π - π stacking arrangement between the aromatic gallate and catechol rings, absent in analogous dimeric procyanidins, which reduces the total surface accessible to **oxidizing** agents.

REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 20 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1996:465752 CAPLUS

DOCUMENT NUMBER: 125:189104

TITLE: Inhibition of β -glucosidase (*Amygdalae dulces*) by (+)-**catechin oxidation** products and procyanidin **dimers**

AUTHOR(S): Guyot, Sylvain; Pellerin, Patrice; Brillouet, Jean-Marc; Cheynier, Veronique

CORPORATE SOURCE: Laboratoire Polymeres Techniques Physico-Chimiques, IPV, Institut National Recherche Agronomique, Montpellier, 34060, Fr.

SOURCE: Bioscience, Biotechnology, and Biochemistry (1996), 60(7), 1131-1135

CODEN: BBBIEJ; ISSN: 0916-8451

PUBLISHER: Japan Society for Bioscience, Biotechnology, and Agrochemistry

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 154-23-4, (+)-**Catechin 23567-23-9**, Procyanidin

B3 29106-49-8, Procyanidin B2 36523-87-2


73086-04-1 179185-59-2 179185-60-5

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study) (inhibition of β -glucosidase from almond by (+)- **catechin oxidation** products and procyanidin **dimers**)

RN 154-23-4 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




10/783,801

RN 23567-23-9 CAPLUS

CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3S,3'S,4S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 29106-49-8 CAPLUS

CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3R,3'R,4R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 36523-87-2 CAPLUS


CN 2H-1-Benzopyran-3,5,7-triol, 8-[2-[(2R,3S)-3,4-dihydro-3,5,7-trihydroxy-2H-

10/783,801


1-benzopyran-2-yl]-4,5-dihydroxyphenyl]-2-(3,4-dihydroxyphenyl)-3,4-dihydro-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A





PAGE 2-A



RN 73086-04-1 CAPLUS

CN Benzoic acid, 3,4,5-trihydroxy-, (2R,2'R,3R,3'R,4R)-2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-3,5,5',7,7'-pentahydroxy[4,8'-bi-2H-1-benzopyran]-3'-yl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 179185-59-2 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 8-[5-(3,4-dihydro-3,5,7-trihydroxy-2H-1-benzopyran-2-yl)-2-hydroxyphenoxy]-2-(3,4-dihydroxyphenyl)-3,4-dihydro-, [2R-[2 α ,3 β ,8(2R*,3S*)]]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 179185-60-5 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 8-[4-(3,4-dihydro-3,5,7-trihydroxy-2H-1-benzopyran-2-yl)-2-hydroxyphenoxy]-2-(3,4-dihydroxyphenyl)-3,4-dihydro-, [2R-[2 α ,3 β ,8(2R*,3S*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




AB The sensitivity and specificity of the inhibition of β -glucosidase from almond (*Amygdalae dulces*) by (+)-**catechin**, an **oxidized** (+)-**catechin** solution, 3 dimeric procyanidins, and 5 (+)-**catechin dimers** obtained by enzymic **oxidation** were evaluated by using a chromatog. **method**. All of the polyphenols tested presented a significant inhibitory effect. Noncompetitive inhibition was observed for the **oxidized** (+)-**catechin** solution. Some **oxidation** products were at least as powerful inhibitors as procyanidins which are known for their tanning effect. Yellow **oxidation** products were among the strongest inhibitors. No marked role of the number of OH and o-diphenol groups nor of the nature or position of the interflavanic linkage in the inhibitory effect was apparent.

L10 ANSWER 21 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN
ACCESSION NUMBER: 1996:420728 CAPLUS

10/783,801



DOCUMENT NUMBER: 125:113062
TITLE: Structural determination of colorless and yellow
dimers resulting from (+)-**catechin**
coupling catalyzed by grape polyphenoloxidase
AUTHOR(S): Guyot, Sylvain; Vercauteren, Joseph; Cheynier,
Veronique
CORPORATE SOURCE: Lab. Polym. Tech. Phys.-Chim., INRA, Montpellier,
34060, Fr.
SOURCE: Phytochemistry (1996), 42(5), 1279-1288
CODEN: PYTCAS; ISSN: 0031-9422
PUBLISHER: Elsevier
DOCUMENT TYPE: Journal
LANGUAGE: English
IT **154-23-4, (+)-Catechin**
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
(Biological study); PROC (Process)
(structure of **dimers** resulting from (+)-**catechin**
oxidation by grape polyphenoloxidase)
RN 154-23-4 CAPLUS
CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-,
(2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



IT **64817-28-3 179185-59-2 179185-60-5**
179185-61-6
RL: BSU (Biological study, unclassified); MFM (Metabolic formation); PRP
(Properties); BIOL (Biological study); FORM (Formation, nonpreparative)
(structure of **dimers** resulting from (+)-**catechin**
oxidation by grape polyphenoloxidase)
RN 64817-28-3 CAPLUS
CN 2H-1-Benzopyran-3,5,7-triol, 8-[6-[(2R,3S)-3,4-dihydro-3,5,7-trihydroxy-2H-
1-benzopyran-2-yl]-2,3-dihydroxyphenyl]-2-(3,4-dihydroxyphenyl)-3,4-
dihydro-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 179185-59-2 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 8-[5-(3,4-dihydro-3,5,7-trihydroxy-2H-1-benzopyran-2-yl)-2-hydroxyphenoxy]-2-(3,4-dihydroxyphenyl)-3,4-dihydro-, [2R-[2 α ,3 β ,8(2R*,3S*)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


10/783,801



RN 179185-60-5 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 8-[4-(3,4-dihydro-3,5,7-trihydroxy-2H-1-benzopyran-2-yl)-2-hydroxyphenoxy]-2-(3,4-dihydroxyphenyl)-3,4-dihydro-, [2R-[2 α ,3 β ,8(2R*,3S*)]]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 179185-61-6 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 6-[2-[(2R,3S)-3,4-dihydro-3,5,7-trihydroxy-2H-1-benzopyran-2-yl]-4,5-dihydroxyphenyl]-2-(3,4-dihydroxyphenyl)-3,4-dihydro-, (2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



AB (+)-**Catechin oxidation** was carried out in aqueous systems using grape polyphenoloxidase as catalyst. Eight fractions corresponding to the major products formed at pH 3 and 6 were purified using HPLC at the **preparative** and semi-**preparative** scale. Structural characterization using UV-visible detection and mass spectrometry indicated that they corresponded to (+)-**catechin dimers** including two yellow pigments. Mono- and bi-dimensional ^1H and ^{13}C NMR analyses provided structural hypotheses for five **oxidation** products whereas the other three fractions were mixts. of several isomers. Colorless products, with C-C or C-O interflavan linkages, were dehydrodicatechins of the B-type. One of the two pigments corresponds to dehydrodicatechin A, already identified in other **oxidation** models, and the other pigment is a new structure of the quinone-methide type.

L10 ANSWER 22 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1996:140282 CAPLUS

DOCUMENT NUMBER: 124:200664

TITLE: Principal phenolic phytochemicals in French Syrah and Grenache Rhone wines and their antioxidant activity in inhibiting **oxidation** of human low density lipoproteins

AUTHOR(S): Teissedre, P.-L.; Waterhouse, A. L.; Frankel, E. N.

CORPORATE SOURCE: Faculte de Pharmacie, Universite de Montpellier I, Montpellier, 34060, Fr.

SOURCE: Journal International des Sciences de la Vigne et du Vin (1995), 29(4), 205-12

CODEN: JISVE8; ISSN: 1151-0285

PUBLISHER: Vigne et Vin Publications Internationales

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 528-58-5, Cyanidin 7228-78-6, Malvidin-3-glucoside

20315-25-7, Procyanidin B1 23567-23-9, Procyanidin B3

29106-49-8, Procyanidin B2


RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence)

(phenolic constituents in French Syrah and Grenache Rhone wines and their antioxidant activities in inhibiting **oxidation** of human low d. lipoproteins)

RN 528-58-5 CAPLUS


CN 1-Benzopyrylium, 2-(3,4-dihydroxyphenyl)-3,5,7-trihydroxy-, chloride (9CI) (CA INDEX NAME)

10/783,801



RN 7228-78-6 CAPLUS
CN 1-Benzopyrylium, 3-(β -D-glucopyranosyloxy)-5,7-dihydroxy-2-(4-hydroxy-3,5-dimethoxyphenyl)-, chloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 20315-25-7 CAPLUS
CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3R,3'S,4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/783,801




RN 23567-23-9 CAPLUS
CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3S,3'S,4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



RN 29106-49-8 CAPLUS
CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3R,3'R,4R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



AB Wine contains natural plant phenolic antioxidants that may protect circulating lipoproteins from oxidative damage. By inhibition of the copper-catalyzed **oxidation** of LDL, we determined the activity of thirteen Rhone Coast wines exclusively from Syrah and Grenache varieties. About 50% of the wines were **made** with a long maceration **process**. Major monomeric phenolic compds. and procyanidin **dimers** were analyzed in each sample by HPLC and correlated with relative LDL antioxidant activity. Correlations obtained can be grouped in 3 classes: **catechin** ($r = 0.75$), procyanidins B1, B2, and B3 ($r = 0.43-0.55$), malvidin-3-glucoside and cyanidin ($r = 0.43$), gallic acid, myricetin, rutin ($r = 0.2-0.4$). On the same basis total phenol contents of wines gave a correlation with LDL antioxidant activity of $r = 0.72$. Comparison, at the same total phenol concentration, with different red California wines shows that antioxidant activity of French Syrah and Grenache range between that of Merlot (56-65%) and Cabernet Sauvignon (37-45%). In contrast Syrah wines **made** with a short extraction **process** gave lower inhibition of LDL **oxidation** of 16% which is less than white California wines averaging 36%. Activity of each wine phenolic compound can play a role in protecting LDL against **oxidation**

L10 ANSWER 23 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1995:792395 CAPLUS

DOCUMENT NUMBER: 123:197162

TITLE: Reactions of enzymically generated quinones in relation to browning in grape musts and wines

AUTHOR(S): Cheynier, Veronique; Fulcrand, Helene; Guyot, Sylvain; Oszmianski, Jan; Moutounet, Michel

CORPORATE SOURCE: Laboratoire des Polymeres et des Technique Physico-chimiques, Institut National de la Recherche Agronomique, Montpellier, 34060, Fr.

SOURCE: ACS Symposium Series (1995), 600(Enzymatic Browning and Its Prevention), 130-43
CODEN: ACSMC8; ISSN: 0097-6156

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 7084-24-4, Cyanidin-3-glucoside


10/783,801

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(reactions of enzymically generated quinones in relation to browning in grape musts and wines)

RN 7084-24-4 CAPLUS

CN 1-Benzopyrylium, 2-(3,4-dihydroxyphenyl)-3-(β -D-glucopyranosyloxy)-5,7-dihydroxy-, chloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 154-23-4, (+)-Catechin 7228-78-6,
Malvidin-3-glucoside

RL: RCT (Reactant); RACT (Reactant or reagent)
(reactions of enzymically generated quinones in relation to browning in grape musts and wines)

RN 154-23-4 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-, (2R,3S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



RN 7228-78-6 CAPLUS

CN 1-Benzopyrylium, 3-(β -D-glucopyranosyloxy)-5,7-dihydroxy-2-(4-hydroxy-3,5-dimethoxyphenyl)-, chloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.



AB The role of enzymically generated caffeoyltartaric acid quinones in browning development was investigated in grape musts and model solns. Caffeoyltartaric acid quinones are only slightly colored and undergo condensation rather slowly, yielding mostly colorless products. On the other hand, they are powerful oxidants and, in particular, **oxidize** flavan-3-ols to unstable secondary quinones which proceed readily to brown polymers. The structures of **dimers** resulting from caffeic acid **oxidation** were elucidated, and the involvement of nucleophilic attack by water in the **process** of their formation demonstrated. Seven **dimers**, including both colorless and brown compds., were obtained by **oxidation** of **catechin** and characterized. The involvement of caffeoyltartaric acid quinone in anthocyanin oxidative degradation was also established. Trapping of primary quinones by glutathione competes with reactions leading to must discoloration but maintains high levels of **oxidizable** compds., especially flavan-3-ols, which serve as browning precursors in wine.

L10 ANSWER 24 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1984:470941 CAPLUS

DOCUMENT NUMBER: 101:70941

TITLE: Turbidity formed in beer at low temperatures.

Affinity of proanthocyanidins and their **oxidation** products for haze-forming proteins of beer and the formation of chill haze

AUTHOR(S): Asano, Katsuhiko; Ohtsu, Keiji; Shinagawa, Kyoko; Hashimoto, Naoki

CORPORATE SOURCE: Brew. Sci. Lab., Kirin Brew. Co., Ltd., Takasaki, 370-12, Japan

SOURCE: Agricultural and Biological Chemistry (1984), 48(5), 1139-46

CODEN: ABCHA6; ISSN: 0002-1369

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 154-23-4 490-46-0 23567-23-9

37064-31-6

RL: PRP (Properties)

(protein affinity of, of beer, chill haze in relation to)


RN 154-23-4 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-,

10/783,801

(2R,3S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



RN 490-46-0 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-,
(2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




RN 23567-23-9 CAPLUS

CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3S,3'S,4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).


10/783,801




RN 37064-31-6 CAPLUS
CN [4,8':4',8''-Ter-2H-1-benzopyran]-3,3',3'',5,5',5'',7,7',7''-nonol,
2,2',2''-tris(3,4-dihydroxyphenyl)-3,3',3'',4,4',4''-hexahydro-,
(2R,2'R,2''R,3S,3'S,3''S,4R,4'R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

PAGE 1-A





AB Among the polyphenols found in beer, proanthocyanidins have a specific affinity for the haze-forming proteins in beer and retained their capacity to form a chill haze. Proanthocyanidin pentamer and tetramer had the highest affinities for the haze-forming proteins, followed by trimer, **dimer**, and **catechins**. During the brewing **process**, proanthocyanidin trimer and polymeric proanthocyanidins easily formed insol. complexes with proteins in the wort as a result of their high affinity for proteins; consequently, these were not found in finished beer, whereas such proanthocyanidin **dimers** as procyanidin B3 [23567-23-9], and **catechin** [154-23-4] survived in finished beer. Procyanidin B3 and **catechin**, when stored in beer or a buffer solution, seemed to undergo oxidative polymerization and increased their affinity for the haze-forming proteins to form a extensive chill haze.

L10 ANSWER 25 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1981:619585 CAPLUS

DOCUMENT NUMBER: 95:219585

TITLE: Stability of (+)-cyanidanol-3 in aqueous solution

AUTHOR(S): Akimoto, Koichi; Sugimoto, Isao

CORPORATE SOURCE: Pharm. Res. Cent., Kanebo Ltd., Osaka, 534, Japan

SOURCE: Chemical & Pharmaceutical Bulletin (1981), 29(7), 2005-11

CODEN: CPBTAL; ISSN: 0009-2363

DOCUMENT TYPE: Journal

LANGUAGE: English


IT 490-46-0P

RL: FORM (Formation, nonpreparative); PREP (Preparation)
(formation of, from cyanidanol, kinetics of)

RN 490-46-0 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-,
(2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



10/783,801


IT 154-23-4

RL: RCT (Reactant); RACT (Reactant or reagent)
(reactions of, in aqueous solution)

RN 154-23-4 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-,
(2R,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



AB Two degradation products were obtained from the title compound (I) at pH 1.4: **epicatechin** (II) and the **dimer**. In the basic pH region, the main product was II. The effect of O was remarkable, especially in basic solution. In strongly acidic solution, the degradation of I occurred by simultaneous

apparent 1st-order (epimerization) and 2nd-order (dimerization) reactions. The rate consts. were calculated by a new **method**. In the neutral and basic pH regions, the reverse reaction between I and II took place and the rate consts. were obtained.

L10 ANSWER 26 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1980:56776 CAPLUS

DOCUMENT NUMBER: 92:56776

TITLE: The procyanidins of white grapes and wines

AUTHOR(S): Lea, Andrew G. H.; Bridle, Peter; Timberlake, Colin F.; Singleton, Vernon L.

CORPORATE SOURCE: Long Ashton Res. Stn., Univ. Bristol, Long Ashton/Bristol, BS18 9AF, UK

SOURCE: American Journal of Enology and Viticulture (1979), 30(4), 289-300

CODEN: AJEVAC; ISSN: 0002-9254

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 4852-22-6D, derivs. 20315-25-7 23567-23-9


29106-49-8 29106-51-2

RL: BIOL (Biological study)
(of grapes and wines)

RN 4852-22-6 CAPLUS


CN 2H-1-Benzopyran-3,4,5,7-tetrol, 2-(3,4-dihydroxyphenyl)-2-[[2-(3,4-dihydroxyphenyl)-3,4-dihydro-5,7-dihydroxy-2H-1-benzopyran-3-yl]oxy]-3,4-dihydro- (9CI) (CA INDEX NAME)

10/783,801



RN 20315-25-7 CAPLUS
CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3R,3'S,4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 23567-23-9 CAPLUS
CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3S,3'S,4S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).

10/783,801




RN 29106-49-8 CAPLUS
CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3R,3'R,4R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 29106-51-2 CAPLUS
CN [4,8'-Bi-2H-1-benzopyran]-3,3',5,5',7,7'-hexol, 2,2'-bis(3,4-dihydroxyphenyl)-3,3',4,4'-tetrahydro-, (2R,2'R,3S,3'R,4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



AB Procyanidin exts. were **prepared** from a Seyval white wine, from a Mueller-Thurgau wine fermented partly on its skins, and from grape seeds. The exts. were studied chromatog. and revealed a range of discrete procyanidin oligomers up to the pentameric and significant quantities of more polymeric and **oxidized** material. Semiquant. estns. showed that the level of total procyanidins was as low as 5-10 mg/L in the Seyval wine rising to 300 mg/L in the Mueller-Thurgau, but that the qual. distribution patterns of the **catechins** and procyanidins showed no marked differences between the two. The Mueller-Thurgau extract was examined counter-current distribution between EtOAc and H₂O, thus achieving a separation of different groups of oligomeric procyanidins, and this was followed by further chromatog. treatment on Sephadex LH-20. The 4 main procyanidin **dimers** of white wines were isolated in the free state and were characterized by chromatog. and their degradative behavior in the presence of acidic toluene-thiol as procyanidins B1-B4. A further procyanidin fraction was isolated and shown to consist of a mixture of 2 stereoisomeric trimers, 1 composed of 3 **epicatechin** units and the other of 2 **epicatechins** and a terminal **catechin**. The English-grown Mueller-Thurgau wine contained negligible amts. of gallocatechins and of galloyl esters of **catechins**, either alone or in combination as procyanidins; this is in contrast to some previous reports.

L10 ANSWER 27 OF 27 CAPLUS COPYRIGHT 2006 ACS on STN

ACCESSION NUMBER: 1965:433098 CAPLUS

DOCUMENT NUMBER: 63:33098

ORIGINAL REFERENCE NO.: 63:5944f-h,5945a

TITLE: Suitability of sulfurous acid for hydrolysis of condensed tannins

AUTHOR(S): Quesnel, V. C.

CORPORATE SOURCE: Univ. West Indies, St. Augustine, Trinidad

SOURCE: Tetrahedron Letters (1964), (48), 3699-702

CODEN: TELEAY; ISSN: 0040-4039

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 490-46-0, Epicatechol


(from Cacao leucocyanidin polymer)

10/783,801

RN 490-46-0 CAPLUS

CN 2H-1-Benzopyran-3,5,7-triol, 2-(3,4-dihydroxyphenyl)-3,4-dihydro-,
(2R,3R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



AB Although condensed tannins are hydrolyzed by dilute acids, concomitant acid-catalyzed polymerization occurs. Alkaline hydrolysis is complicated by oxidative condensation. The suitability of H_2SO_3 , as suggested by Forsyth and Roberts (CA 54, 14238b) in a study of the hydrolysis of leucocyanidin **dimer**, for investigations on the structure of condensed tannins was examined. Acid-catalyzed polymers were **prepared** according to the **method** of Goldstein and Swain (CA 60, 2046d) and the precipitated polymers collected by filtration. **Oxidation** polymers were **prepared** from 100-mg. samples in 50 ml. H_2O by overnight **oxidation** at 20° with 75 mg. NaIO_4 ; the solns. were saturated with salt and the polymers collected on sintered glass. Cacao polyphenyl storage cells (Brown, Nature 173(4402), 492(1954)) and mangrove bark, extracted with MeOH and the exts. diluted with 2 vols. Et_2O , gave cacao and mangrove tannins, resp. Polymerization by acid and by **oxidation** gave (-)-**epicatechin** polymers (I, II) and cacao leucocyanidins 1 and 2 polymers (III, IV). Extraction gave polymeric cacao leucocyanidin (V, VI) from cacao and mangrove tannin, resp. I and II were practically insol. in H_2O ; the others were H_2O -soluble. I was partially and III wholly soluble in EtOH. On 2-way chromatography in 2% aqueous AcOH and a BuOH phase of 4:1:5 BuOH-AcOH- H_2O , none showed any discreet spot in both solvents. The polymers were refluxed in aqueous or alc. H_2SO_3 for 2.5 hrs. and again examined chromatographically. No spot mobile in both solvents was produced from I, III, or IV, but the others gave mobile spots. II was split to give a spot with R_f 0.02 (2% aqueous HOAc), 0.5 (BuOH-HOAc- H_2O , 4:1:5) and another with R_f 0.95 (aqueous HOAc), 0.5 (BuOH). V was completely split into (-)-**epicatechin**, substance A, and substance B. Longer treatment gave complete hydrolysis to substance A, (-)-**epicatechin**, and (+)-**catechin**. Hydrolysis of VI was extensive. The bulk of VI was closely similar to that of V and it is possible that other condensed tannins are similar to V. The use of H_2SO_3 in their structural investigations was suggested. Quebracho tannin is extensively altered by bisulfite treatment.

=> log y

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

180.54

347.69

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

10/783,801

	ENTRY	SESSION
CA SUBSCRIBER PRICE	-20.25	-20.25

STN INTERNATIONAL LOGOFF AT 09:46:53 ON 07 JUN 2006